University of Maryland





 $\mathscr{R}(D^{(*)})$ ,  $\mathscr{R}_{K^{(*)}}$ , and their cousins: update on the continued challenges to lepton flavor universality

### 5<sup>th</sup> May 2021

Virtual joint JHU/UMD seminar











### ~Assume Universe is $SU(3) \times SU(2)_L \times U(1)$ symmetric, put in a few particles, and bam!, most precise and comprehensive theory in the history of mankind



Manuel Franco Sevilla

# The SM is pretty good



Anomalous magnetic dipole moment



12,672 diagrams of 10<sup>th</sup> order





















## Beyond the SM



#### Looking in many directions

















By Pallab Ghosh Science corresponder

O 7 Apri















#### Manuel Franco Sevilla

## Testing the SM





#### Machine finds tantalising hints of



### ~Alas, **no direct** detection yet

Can access mass scales beyond the reach of current particle accelerators through **precision** 

### tests

→ Flavor physics (study of quark and lepton species) is a key tool

Slide 4



## Testing the SM



#### Manuel Franco Sevilla



### ~Alas, **no direct** detection yet

Can access mass scales beyond the reach of current particle accelerators through precision

### tests

→ Flavor physics (study of quark and lepton species) is a key tool

Slide 4



# Lepton Flavor Universality (LFU)

### ∼ It is assumed that electroweak gauge couplings to 3 fermion generations are identical





Manuel Franco Sevilla











## LFU tested to great precision

#### LFU tests with 1<sup>st</sup>/2<sup>nd</sup> gen.

To **0.28%** in Z decays

> To **0.8%** in W decays

To **0.31%** in meson decays

To 0.14% in  $\tau \rightarrow \ell \nu \nu$ 

$\Gamma_{Z \to \mu\mu}$	1 0000 - 0 0028
$\Gamma_{Z \to ee}$	LEP, <u>Phys. Rept. 427 (2006) 257</u>
$\mathscr{B}(W \to \phi)$ $\mathscr{B}(W \to \rho)$	$(2\nu)$ $(2\nu)$ = 1.004 ± 0.008 (2019) CDF + LHC, JPG: NPP, 46, 2 (2019)
$\frac{\Gamma_{J/\psi \to \mu\mu}}{\Gamma} =$	$= 1.0016 \pm 0.0031$
	DG (BESIII), <u>RPP, Chin. Phys. C40 (2016) 100001</u> ( $2.488 \pm 0.009$ ) × $10^{-5}$
$\Gamma_{K \to \mu\nu}$ $\Gamma_{\pi \to e\nu}$	PDG (NA62), <u>RPP, Chin. Phys. C40 (2016) 100001</u> (1 230 + 0 004) $\times$ 10 <sup>-4</sup>
$\Gamma_{\pi \to \mu \nu}$	PiENu, <u>Phys. Rev. Lett. 115, 071801 (2015</u> )

 $g_{\mu}/g_e = 1.0018 \pm 0.0014$ PDG, A. Pich, Prog. Part. Nucl. Phys. 75 (2014) 41

Manuel Franco Sevilla



#### LFU tests with 3<sup>rd</sup> gen.

To **0.32%** in Z decays

**2.6** $\sigma$  tension in W decays

> To 1.3% in W decays

 $\frac{\Gamma_{Z \to \tau \tau}}{1} = 1.0019 \pm 0.0032$  $\Gamma_{Z \rightarrow ee}$ LEP, Phys. Rept. 427 (2006) 257  $\frac{\Gamma_{W \to \tau \nu}}{1.070 \pm 0.026}$  $\Gamma_{W \to \mu \nu}$ LEP, Phys. Rept. 532 (2013) 119,  $\frac{\Gamma_{W \to \tau \nu}}{1} = 0.992 \pm 0.013$ 

 $\Gamma_{W \to \mu \nu}$ ATLAS, arXiv:2007.14040

To **6.1%** in  $D_{s}$  decays

 $\frac{\Gamma_{D_s \to \tau \nu}}{1} = 9.95 \pm 0.61$  $\Gamma_{D_s \to \mu\nu}$ HFLAV, Eur. Phys. J. C77 (2017) 895

To **0.15%** in  $\tau \rightarrow \ell \nu \nu \text{ (with } \tau_{\tau})$ 

 $g_{\tau}/g_{\mu} = 1.0030 \pm 0.0015$ 

PDG, S. Pich, Prog. Part. Nucl. Phys. 75 (2014) 41



### ~ Since 2012, hints of LFU in transitions involving 3<sup>rd</sup> gen. b quark



Very solid SM predictions with 1-2% uncertainty, established deviations would be clear indications of BSM physics

Manuel Franco Sevilla

 $\mathscr{R}(D^{(*)})$ ,  $\mathscr{R}_{K^{(*)}}$ , and their cousins: update on the continued challenges to LFU

## B anomalies



Slide 7





## Overview of experiments

### LFU results with $b \rightarrow c \tau \nu$

- $p_R$  reconstruction
- B-factory and LHCb measurements of  $\mathscr{R}\left(D^{(*)}
  ight)$
- Beyond  $\mathscr{R}(D^{(*)})$
- Future prospects

## Outline





## **One elegant** interpretation









## Contributions from several experiments 🖓 👘



Manuel Franco Sevilla



 $\mathcal{O}(10^9) B^{0/+}$  mesons Low uncertainty on absolute rates, 100% ε(trigger), PID, low e-brem, knowledge of collision momentum

### **B-factories**



With  $\mathcal{O}(10^8) B^{0/+}$  mesons already competitive search for  $B \to K \bar{\nu} \bar{\nu}$  (backup)!

 $\mathcal{O}(10^{11}) B_{(s)}^{0/+}$  mesons Triggers primarily for flavor, PID, VELO, all b-hadron species

LHC

 $\mathcal{O}(10^{12}) B_{(c)}^{0/+}$  mesons **All b-hadron species** 







# LHC environment is slightly busier

 $pp \to X_b B_s^0 X$  $B_s^0 \rightarrow \mu^+ \mu^-$ 



**LHC** pp collisions have background from  $b\bar{b}$  hadronization, underlying event, and pileup

> Clean  $e^+e^-$  collisions only produce two B mesons (for the most part)

Manuel Franco Sevilla



### **B-factories**







# Vertexing and isolation key to LHC



Manuel Franco Sevilla



~ B mesons can fly ~ cm thanks to large boost ~ Excellent trackers in CMS and ATLAS ~ Superb vertexing by VELO in LHCb → Only 8.2 mm from IP, reduced to 5.1 in upgrade Multivariate algorithms ensure tracks isolated Based on track impact parameter, other variables











# LFU results with $b \rightarrow c \tau \nu$ transitions

Driginally from BaBa





### Leptonic τ

 $\tau^- \to \ell^- \nu_\tau \bar{\nu}_\ell$ 

Same reco particles as normalization  $B \rightarrow D^{(*)} \ell \nu$ , many uncertainties cancel on  $\mathscr{R}\left(D^{(*)}\right)$ 



$$\mathscr{R}\left(D^{(*)}\right) = \frac{\mathscr{B}\left(\bar{B} \to D^{(*)}\tau\nu_{\tau}\right)}{\mathscr{B}\left(\bar{B} \to D^{(*)}\ell\nu_{\ell}\right)} = \frac{N_{sig}}{N_{norm}}\frac{\epsilon_{norm}}{\epsilon_{sig}}$$

ε ratio easy, yields are key

#### Manuel Franco Sevilla

## τ reconstruction







# $p_R$ reconstruction at the B-factories

**B tagging:**  $p_{B_{sig}} = p_{e^+e^-} - p_{B_{tag}}$ <u>Hadronic</u>: best  $\sigma(p_R)$ ,  $\epsilon_{had} \sim 0.2-0.4\%$ <u>Semileptonic</u>: worse  $\sigma(p_R)$ ,  $\epsilon_{sl} \sim 0.3-0.6\%$  $B^-$ ,  $ar{B}^0$  ( FEI: bottom-up approach based on BDTs with  $\epsilon_{FEI}$  up to 3x the corresponding  $\epsilon_{had}$  or  $\epsilon_{sI}$ 

















# Belle 2015 $\mathscr{R}(D^{(*)})$ , q<sup>2</sup> distributions



Phys. Rev. D **92**, 072014 (2015)

Similar strategy to BaBar

 $\Rightarrow$  Hadronic  $B_{tag}$ , leptonic  $\tau$ 

 $\sim$  Also excess, consistent with BaBar



Manuel Franco Sevilla



# $p_R$ reconstruction at LHCb



 $\mathscr{R}(D^{(*)})$ ,  $\mathscr{R}_{K^{(*)}}$ , and their cousins: update on the continued challenges to LFU

Manuel Franco Sevilla



Slide 17







![](_page_19_Picture_4.jpeg)

![](_page_20_Picture_0.jpeg)

# Beyond $\mathscr{R}(D^{(*)})$

- ~ Even a 5 $\sigma$  on  $\mathscr{R}(D^{(*)})$  would not be sufficient to convince ourselves of NP
  - Indirect measurement with broad signal distributions due to multiple v in final state
- It will be important to have
  - Confirmation by independent experiments
  - Confirmation in different decays
  - Characterization in kinematic distributions

Belle II and upgraded LHCb both **sensitive** to angular distributions

> Hill, John, Ke, Poluektov, *JHEP* 2019, 133 (2019)

![](_page_20_Figure_10.jpeg)

Manuel Franco Sevilla

![](_page_20_Picture_12.jpeg)

LHCb has a unique ability to study  $b \rightarrow c \tau \nu$ transitions because bb production at the LHC hadronizes into all species of b-hadrons

![](_page_20_Figure_16.jpeg)

![](_page_20_Figure_17.jpeg)

![](_page_20_Figure_18.jpeg)

LHCb already published first non- $\mathscr{R}(D^{(*)})$  measurement  $\mathscr{R}(J/\Psi) = 0.71 \pm 0.17 \pm 0.18,$ **1.8σ above SM** 

Phys. Rev. Lett. 120, 121801 (2018)

![](_page_20_Picture_21.jpeg)

![](_page_20_Figure_24.jpeg)

![](_page_20_Picture_25.jpeg)

![](_page_21_Picture_0.jpeg)

# Future prospects for LFU in $b \rightarrow c \tau \nu$

~ Currently, world-averaged  $\mathscr{R}(D^{(*)})$  exceeds SM by ~14%

- ~With Belle II and upgraded LHCb, could get uncertainties below 3% in a few years In addition to  $\mathscr{R}(D^{(*)})$  and  $\mathscr{R}(J/\Psi)$ , LHCb has  $\mathscr{R}(D^{**})$ ,  $\mathscr{R}(p\bar{p})$ ,  $\mathscr{R}(D_s)$ ,  $\mathscr{R}(D_s)$ , and  $\mathscr{R}(\Lambda_c)$  ongoing! Even CMS trying to get out a measurement with ingenious trigger strategy

![](_page_21_Figure_6.jpeg)

Manuel Franco Sevilla

![](_page_21_Picture_8.jpeg)

Data sample up to year

Wherever this ends up, very exciting times ahead!

Bernlochner, MFS, Robinson, Wormser, arXiv:2101.08326

![](_page_21_Picture_18.jpeg)

![](_page_21_Picture_20.jpeg)

![](_page_22_Picture_0.jpeg)

Penguin from Jeff Brass

# LFU results with $b \rightarrow see$ transitions

![](_page_23_Figure_0.jpeg)

![](_page_24_Figure_0.jpeg)

![](_page_24_Picture_4.jpeg)

![](_page_25_Figure_0.jpeg)

![](_page_25_Picture_4.jpeg)

![](_page_26_Picture_0.jpeg)

### Not as suppressed as leptonic decays, but still rare with $\mathscr{B} \sim 10^{-7}$

![](_page_26_Figure_3.jpeg)

Manuel Franco Sevilla

## Semileptonic $B_{(s)} \rightarrow H\ell^+\ell^-$ (medium rare)

![](_page_26_Figure_7.jpeg)

![](_page_26_Picture_10.jpeg)

Slide 26

## Precision test strategies

![](_page_27_Picture_1.jpeg)

### ~ Experimental and theoretical uncertainties depend on strategy

![](_page_27_Figure_3.jpeg)

#### **Branching fractions**

Simpler for LHC (focus on  $\mu$ ), but large theory uncertainties

Angular observables Minimal FF uncertainties, though sensitive to charm loops

Manuel Franco Sevilla

![](_page_27_Picture_8.jpeg)

![](_page_27_Figure_9.jpeg)

<u>**LFU ratios</u>**  $\mathscr{R}_{H_s} = \frac{\mathscr{B}(H_b \to H_s \mu \mu)}{\mathscr{B}(H_b \to H_s ee)}$ </u>

Theory uncertainty of ~1%, but electrons harder at the LHC

![](_page_27_Picture_15.jpeg)

![](_page_27_Picture_16.jpeg)

## Differential BF rates

![](_page_28_Picture_1.jpeg)

### ~ First measurements of $B \to K^{(*)} \ell \ell$ at Tevatron and the B-factories - Consistent with expectations though large uncertainties

![](_page_28_Figure_4.jpeg)

.9 Martuel Franco Sevilla m<sub>κπ</sub> (GeV/c<sup>2</sup>)

 $\mathscr{R}(D^{(*)})$ ,  $\mathscr{R}_{K^{(*)}}$ , and their cousins: update on the continued challenges to LFU

![](_page_28_Picture_7.jpeg)

#### **Deficit** in LHCb measurements with muons at low q<sup>2</sup>

![](_page_28_Picture_10.jpeg)

![](_page_28_Picture_11.jpeg)

![](_page_28_Picture_12.jpeg)

![](_page_28_Picture_13.jpeg)

![](_page_29_Figure_0.jpeg)

![](_page_30_Picture_0.jpeg)

![](_page_30_Figure_1.jpeg)

## $P'_5$ and $Q_{4,5}$ in $B \to K^* \ell \ell$

![](_page_30_Picture_4.jpeg)

![](_page_30_Picture_7.jpeg)

![](_page_30_Picture_8.jpeg)

![](_page_31_Picture_0.jpeg)

![](_page_31_Picture_1.jpeg)

$$\sim \text{Measured all isospin variants for } \mathscr{R}_{K^{(*)}} = \frac{\mathscr{B}(B \to K^{(*)}\mu\mu)}{\mathscr{B}(B \to K^{(*)}ee)}$$

$$\sim \text{Fit } M_{bc} = \sqrt{E_{\text{beam}}^2 - p_B^2}$$

$$\quad \cdot \mathscr{R}_{K} \text{ also fits NN and } \Delta E = E_B - E_{\text{beam}}, \mathscr{R}_{K^*} \text{ cuts on them}$$

$$\quad \sim \text{Similar mass resolution for } \mu \text{ and } e$$

$$\quad \cdot \text{Powerful check with } B \to J/\psi(\to \ell\ell) K^{(*)}$$

$$r_{J/\psi}^{K} = \frac{\mathscr{R}[B \to K J/\psi(\to \mu\mu)]}{\mathscr{R}[B \to K J/\psi(\to ee)]} = 0.994 \pm 0.015 \quad r_{J/\psi}^{K^*} = \frac{\mathscr{R}[B \to K^* J/\psi(\to \mu\mu)]}{\mathscr{R}[B \to K^* J/\psi(\to ee)]} = 1.015 \pm 0.045$$

$$\frac{\varphi^{0^{0^{-1}}}_{gam}} = \frac{\mathscr{R}(B^+ \to J/\psi(K^+) = (1.032 \pm 0.025) \times 10^{-3}}{\mathscr{R}(B \to J/\psi) K^*}$$

![](_page_31_Picture_6.jpeg)

![](_page_31_Picture_9.jpeg)

![](_page_32_Figure_0.jpeg)

![](_page_33_Figure_0.jpeg)

![](_page_34_Picture_0.jpeg)

# LFU $\mathscr{R}_{K^{(*)}}$ at LHCb: bkgs & signal shape $\overset{\sim}{\gg}$

![](_page_34_Figure_2.jpeg)

Manuel Franco Sevilla

![](_page_34_Picture_5.jpeg)

![](_page_34_Picture_6.jpeg)

## nds reduced with

- variant masses, eg  $m(K^+e) > m(D^0)$ classifiers
- orial and partially-reco bkgs free in fit

# $\sim B \rightarrow K^{(*)} J/\psi(\rightarrow \ell\ell)$ contamination from

~ Signal shapes taken from simulation

Small corrections obtained from clean  $B \to K^{(*)}J/\psi(\to \ell\ell)$ 

Slide 34

![](_page_35_Figure_0.jpeg)

 $\mathscr{R}(D^{(*)})$ ,  $\mathscr{R}_{K^{(*)}}$ , and their cousins: update on the continued challenges to LFU

Slide 35

![](_page_36_Picture_0.jpeg)

![](_page_36_Picture_1.jpeg)

![](_page_36_Figure_2.jpeg)

![](_page_36_Figure_4.jpeg)

![](_page_37_Figure_0.jpeg)

![](_page_37_Picture_9.jpeg)

# One elegant interpretation

![](_page_38_Picture_1.jpeg)

Based on Isidori at APS April 2021 and Cornella, Faroughy, Fuentes-Martín, Isidori, Neubert - arXiv:2103.16558

![](_page_38_Picture_3.jpeg)

Penguin from Jeff Brassard

![](_page_38_Picture_5.jpeg)

![](_page_39_Picture_0.jpeg)

![](_page_39_Picture_1.jpeg)

![](_page_39_Figure_2.jpeg)

- 1.8 $\sigma$  excess in  $\mathcal{R}(J/\Psi)$
- 14% excess in  $\mathscr{R}(D)$
- 14% excess in  $\mathscr{R}(D^*)$

![](_page_39_Picture_6.jpeg)

![](_page_39_Figure_7.jpeg)

![](_page_39_Picture_11.jpeg)

## LFU results with $b \rightarrow s \ell \ell$

- **Deficit** in differential BF rates with  $\mu$
- 22% deficit in  $B_{(s)}^0 \rightarrow \mu^+ \mu^-$
- 15% deficit in  $\mathcal{R}_{K}$
- ~30% **deficit** in  $\mathcal{R}_{K^*}$

![](_page_39_Picture_17.jpeg)

- Disagreement in  $B \rightarrow K^* \ell \ell$  angular  $P'_5$ 

![](_page_39_Figure_19.jpeg)

![](_page_39_Picture_22.jpeg)

![](_page_39_Picture_23.jpeg)

![](_page_40_Picture_0.jpeg)

### ~ **Dimension-6 operators** identified as relevant set for combined explanation of both anomalies

$$\mathcal{L}_{b \to s\,\ell^+\ell^-} = \frac{4G_F}{\sqrt{2}} \, V_{ts}^* V_{tb} \, \sum_i \, \mathcal{C}_i^\ell \, \mathcal{O}_i^\ell$$

![](_page_40_Picture_4.jpeg)

**K**(\*) B

**Easy and "clean"** 

*Four-quark operators:*  $\mathcal{O}_2 = (\bar{s}_L \gamma_\mu b_L) (\bar{c}_L \gamma_\mu c_L)$ 

![](_page_40_Picture_8.jpeg)

Manuel Franco Sevilla

 $\mathsf{EFT} \ \mathsf{for} \ b \to \mathcal{SCC}$ 

![](_page_40_Picture_12.jpeg)

$$\mathcal{O}_{LL}^{ij\alpha\beta} = (\bar{q}_L^i \gamma_\mu \ell_L^\alpha) (\bar{\ell}_L^\beta \gamma^\mu q_L^j)$$

 $\mathcal{O}_{LR}^{ij\alpha\beta} = (\bar{q}_L^i \gamma_\mu \ell_L^\alpha) (\bar{e}_R^\beta \gamma^\mu d_R^j)$ 

 $\mathcal{O}_{RR}^{ij\alpha\beta} = (\bar{d}_R^i \gamma_\mu e_R^\alpha) (\bar{e}_R^\beta \gamma^\mu d_R^j)$ 

arXiv:2103.16558

#### Separate NP contributions between **Lepton Flavor Universal**

$$\Delta C_{9,10}^U \equiv C_{9,10}^e - C_{9,10}^{SM}$$

and LFU-breaking  $\Delta C_{9,10}^{\mu} \equiv C_{9,10}^{\mu} - C_{9,10}^{e} = C_{9,10}^{\mu} - (C_{9,10}^{SM} + \Delta C_{9,10}^{U})$ **Difficult and** Induce  $\Delta C_{0}^{U}$  but no LFU breaking terms

•  $(\mathscr{R}_{K^{(*)}})$  or axial-current contributions  $(B_s^0 \rightarrow \mu^+ \mu^-)$ 

![](_page_40_Picture_22.jpeg)

![](_page_40_Picture_23.jpeg)

![](_page_40_Picture_24.jpeg)

![](_page_40_Picture_26.jpeg)

![](_page_41_Picture_0.jpeg)

![](_page_41_Figure_2.jpeg)

 $\mathscr{R}(D^{(*)})$ ,  $\mathscr{R}_{K^{(*)}}$ , and their cousins: update on the continued challenges to LFU

EFT fits for  $b \rightarrow s \ell \ell$ 

![](_page_41_Picture_6.jpeg)

scaling with fermion generation

Isidori at APS April 2021,

![](_page_41_Picture_10.jpeg)

![](_page_41_Picture_11.jpeg)

![](_page_42_Picture_0.jpeg)

# remarkably consistent results

![](_page_42_Figure_3.jpeg)

#### Manuel Franco Sevilla

## $b \rightarrow s\ell\ell$ and $b \rightarrow c\tau\nu$

![](_page_42_Figure_6.jpeg)

Isidori at APS April 2021,

![](_page_42_Picture_13.jpeg)

![](_page_42_Picture_14.jpeg)

![](_page_43_Picture_0.jpeg)

# U<sub>1</sub> leptoquark fits all low-energy data

- Gripaios, '10 Marzocca '18

resonances

![](_page_43_Figure_10.jpeg)

#### Manuel Franco Sevilla

![](_page_43_Picture_13.jpeg)

![](_page_44_Picture_0.jpeg)

# U<sub>1</sub> leptoquark within reach

![](_page_44_Figure_2.jpeg)

#### Manuel Franco Sevilla

![](_page_44_Picture_5.jpeg)

Direct LQ searches at LHC have limited mass reach, but **high p**<sub>T</sub> tails in  $\tau\tau$  events would have **sensitivity** at HL-LHC

Also,  $b \rightarrow d\mu\mu$ ,  $b \rightarrow s\tau\tau$ ,  $b \rightarrow s \tau \mu$ , B<sub>s</sub> mixing,  $b \rightarrow s \nu \nu, \tau \rightarrow \mu \mu \mu$ 

Isidori at APS April 2021,

![](_page_44_Picture_11.jpeg)

![](_page_44_Picture_12.jpeg)

![](_page_44_Picture_13.jpeg)

![](_page_44_Picture_14.jpeg)

![](_page_45_Picture_0.jpeg)

![](_page_45_Picture_1.jpeg)

- ~Excesses in decays involving  $b \rightarrow c \tau \nu$  transitions  $\Rightarrow$  3.1 $\sigma$  significance
- ~ Deficits in decays involving  $b \rightarrow s\mu\mu$  transitions
  - → At least  $3.9\sigma$  significant
- ~ U<sub>1</sub> leptoquark could explain both → Within reach at HL-LHC
- Exciting times ahead
  - → LHC still analyzing Runs 1+2 data
  - → Run 3 to start next year with 5x inst. lumi at LHCb
  - Belle II will increase B-factories dataset by 50x
  - → HL-LHC will increase current dataset by 100x

## Conclusions

![](_page_45_Picture_14.jpeg)

#### 18 16 uncertainty [%] 1410 Total ı Optimistic Data sample up to year 0.25tainty 0.50 Ŭ 0.15 ed Projecte 0.050.002025 2020 2015

Run 2

Run 3

Run 4 Run 5

![](_page_45_Figure_16.jpeg)

 $\mathscr{R}(D^{(*)})$ ,  $\mathscr{R}_{K^{(*)}}$ , and their cousins: update on the continued challenges to LFU

2030

Year

![](_page_45_Figure_20.jpeg)

![](_page_45_Figure_21.jpeg)

![](_page_45_Picture_22.jpeg)